



# Brief Suicide Safety Assessment

## Ask **Suicide-Screening** Questions

**What to do when a pediatric patient screens positive for suicide risk:**

- Use after a patient (**10 - 24 years**) screens positive for suicide risk on the asQ
- Assessment guide for mental health clinicians, MDs, NPs, or PAs
- Prompts help determine disposition

### 1 Praise patient *for discussing their thoughts*

“I’m here to follow up on your responses to the suicide risk screening questions. These are hard things to talk about. Thank you for telling us. I need to ask you a few more questions.”

### 2 Assess the patient *(If possible, assess patient alone depending on developmental considerations and parent willingness.)*

Review patient’s responses from the asQ

#### Frequency of suicidal thoughts

Determine if and how often the patient is having suicidal thoughts.

**Ask the patient:** “In the past few weeks, have you been thinking about killing yourself?” **If yes, ask:** “How often?” (once or twice a day, several times a day, a couple times a week, etc.) “When was the last time you had these thoughts?”

“Are you having thoughts of killing yourself right now?” **(If “yes,” patient is at imminent risk and requires an urgent/STAT mental health evaluation and cannot be left alone. Notify patient’s medical team.)**

#### Suicide plan

Assess if the patient has a suicide plan, regardless of how they responded to any other questions (ask about method and access to means).

**Ask the patient:** “Do you have a plan to kill yourself?” **If yes, ask:** “What is your plan?” **If no plan, ask:** “If you were going to kill yourself, how would you do it?”

**Note: If the patient has a very detailed plan, this is more concerning** than if they haven’t thought it through in great detail. If the plan is feasible (e.g., if they are planning to use pills and have access to pills), this is a reason for greater concern and removing or securing dangerous items (medications, guns, ropes, etc.).

#### Past behavior

Evaluate past self-injury and history of suicide attempts (method, estimated date, intent).

**Ask the patient:** “Have you ever tried to hurt yourself?” “Have you ever tried to kill yourself?”

**If yes, ask:** “How? When? Why?” and assess intent: “Did you think [method] would kill you?” “Did you want to die?” (for youth, intent is as important as lethality of method) **Ask:** “Did you receive medical/psychiatric treatment?”

**Note: Past suicidal behavior is the strongest risk factor for future attempts.**

#### Symptoms **Ask the patient about:**

**Depression:** “In the past few weeks, have you felt so sad or depressed that it makes it hard to do the things you would like to do?”

**Anxiety:** “In the past few weeks, have you felt so worried that it makes it hard to do the things you would like to do or that you feel constantly agitated/on-edge?”

**Impulsivity/Recklessness:** “Do you often act without thinking?”

**Hopelessness:** “In the past few weeks, have you felt hopeless, like things would never get better?”

**Anhedonia:** “In the past few weeks, have you felt like you couldn’t enjoy the things that usually make you happy?”

**Isolation:** “Have you been keeping to yourself more than usual?”

**Irritability:** “In the past few weeks, have you been feeling more irritable or grouchy than usual?”

**Substance and alcohol use:** “In the past few weeks, have you used drugs or alcohol?” **If yes, ask:** “What? How much?”

**Sleep pattern:** “In the past few weeks, have you had trouble falling asleep or found yourself waking up in the middle of the night or earlier than usual in the morning?”

**Appetite:** “In the past few weeks, have you noticed changes in your appetite? Have you been less hungry or more hungry than usual?”

**Other concerns:** “Recently, have there been any concerning changes in how you are thinking or feeling?”

#### Social Support & Stressors

*(For all questions below, if patient answers yes, ask them to describe.)*

**Support network:** “Is there a trusted adult you can talk to? Who? Have you ever seen a therapist/counselor?” **If yes, ask:** “When?”

**Family situation:** “Are there any conflicts at home that are hard to handle?”

**School functioning:** “Do you ever feel so much pressure at school (academic or social) that you can’t take it anymore?”

**Bullying:** “Are you being bullied or picked on?”

**Suicide contagion:** “Do you know anyone who has killed themselves or tried to kill themselves?”

**Reasons for living:** “What are some of the reasons you would NOT kill yourself?”

Ask **Suicide-Screening** Questions**3 Interview patient & parent/guardian together**

If patient is  $\geq 18$  years, ask patient's permission for parent/guardian to join.

**Say to the parent:** "After speaking with your child, I have some concerns about his/her safety. We are glad your child spoke up as this can be a difficult topic to talk about. We would now like to get your perspective."

- "Your child said... (reference positive responses on the asQ). Is this something he/she shared with you?"
- "Does your child have a history of suicidal thoughts or behavior that you're aware of?" **If yes, say:** "Please explain."
- "Does your child seem:
  - o Sad or depressed?"
  - o Anxious?"
  - o Impulsive? Reckless?"
  - o Hopeless?"
  - o Irritable?"
  - o Unable to enjoy the things that usually bring him/her pleasure?"
  - o Withdrawn from friends or to be keeping to him/herself?"

- "Have you noticed changes in your child's:
  - o Sleeping pattern?"
  - o Appetite?"
- "Does your child use drugs or alcohol?"
- "Has anyone in your family/close friend network ever tried to kill themselves?"
- "How are potentially dangerous items stored in your home?" (e.g. guns, medications, poisons, etc.)
- "Does your child have a trusted adult they can talk to?" (Normalize that youth are often more comfortable talking to adults who are not their parents)
- "Are you comfortable keeping your child safe at home?"

**At the end of the interview, ask the parent/guardian:** "Is there anything you would like to tell me in private?"

**4 Make a safety plan with the patient** Include the parent/guardian, if possible.

Create a safety plan for managing potential future suicidal thoughts. A safety plan is different than making a "safety contract"; asking the patient to contract for safety is NOT effective and may be dangerous or give a false sense of security.

**Say to patient:** "Our first priority is keeping you safe. Let's work together to develop a safety plan for when you are having thoughts of suicide."

Examples: "I will tell my mom/coach/teacher."  
"I will call the hotline." "I will call \_\_\_\_\_."

Discuss coping strategies to manage stress (such as journal writing, distraction, exercise, self-soothing techniques).

**Discuss means restriction** (securing or removing lethal means): "Research has shown that limiting access to dangerous objects saves lives. How will you secure or remove these potentially dangerous items (guns, medications, ropes, etc.)?"

**Ask safety question:** "Do you think you need help to keep yourself safe?" (A "no" response does not indicate that the patient is safe; but a "yes" is a reason to act immediately to ensure safety.)

**5 Determine disposition**

After completing the assessment, choose the appropriate disposition plan.

- Emergency psychiatric evaluation:** Patient is at imminent risk for suicide (current suicidal thoughts). Keep patient safe on the unit. Follow the standard of care for a suicidal patient (e.g. remove dangerous objects, 1:1 observer). Request a STAT, emergency psychiatric evaluation.
- Further evaluation of risk is necessary:** Request a comprehensive mental health/safety evaluation prior to discharge.
- Patient might benefit from non-urgent mental health follow-up post-discharge:** No further mental health evaluation in the hospital is needed at this time. Review safety plan for potential future suicidal thoughts and refer patient for a follow-up mental health evaluation in the community, post-discharge.
- No further intervention is necessary at this time.**

**6 Provide resources to all patients**

- 24/7 National Suicide Prevention Lifeline 1-800-273-TALK (8255) En Español: 1-888-628-9454
- 24/7 Crisis Text Line: Text "HOME" to 741-741

